


CRYSTAL


Deep





CRYSTAL

Ultra



CRYSTAL®

CRYSTAL Lite

CRYSTAL Deep

CRYSTAL Market



CRYSTAL Fillers

A line of premium fillers that are created using NASHA technology – an advanced process of purifying hyaluronic acid (HA) without using any animal-sourced materials, ensuring that the risk of impurities is close to zero.

The fillers also contain lidocaine for reducing pain and discomfort.

CRYSTAL fillers are suitable for correcting wrinkles, reducing the depth of nasolabial folds and folds in the corners of the mouth (puppet lines), and are widely used for correcting aesthetic defects of minor tissue deficiencies.

The line consists of:

- Crystal Lite
- Crystal Deep
- Crystal Ultra





CRYSTAL Lite

Correction of fine lines and minimal aesthetic defects

Depth of introduction:

• Intradermally (surface to mid dermis)

Indications for use:

- Elimination of irregularities of the skin, including the back of the hands (thin skin)
- Correction of superficial wrinkles ("pouch", "crow's feet", wrinkles in the frontal and inner brow areas (I-II degree of severity), preauricular wrinkles)
- Correction of nasolacrimal and palpebromalar lines
- Correction of atrophic scars
- Contouring of the face, neck and décolleté



CRYSTAL Lite

Product Details

PACKAGE	1 x 1.0 ml
HA CONCENTRATION	20 mg/ml
LIDOCAINE	0.3 %
NEEDLES IN ASSORTMENT	27 G ½" – 1 ea. 30 G ½" – 1 ea.
STABILIZATION	BDDE
CROSS-LINK LEVEL	•
DURATION*	6-12 months

Package

1 box: 1 syringe / 2 needles

* Duration period can depend on the injection method, area of injection and the patient's individual characteristics.



Storage

Avoid direct sunlight and store at 2°C to 25°C



CRYSTAL Deep

Correction of prominent lines and deep wrinkles



• Intradermally (mid to deep dermis)

Indications for use:

- Correction of aesthetic defects
- Elimination of wrinkles and folds of medium depth: nasolabial, labial wrinkles
- Correction of age-related hand changes
- Correction of earlobes
- Volumetric lip correction



CRYSTAL Deep

Product Details

PACKAGE	1 x 1.0 ml
HA CONCENTRATION	20 mg/ml
LIDOCAINE	0.3 %
NEEDLES IN ASSORTMENT	27G ½" - 2 ea.
STABILIZATION	BDDE
CROSS-LINK LEVEL	
DURATION*	9-12 months

Package

1 box: 1 syringe / 2 needles

* Duration period can depend on the injection method, area of injection and the patient's individual characteristics.



Storage

Avoid direct sunlight and store at 2°C to 25°C

CRYSTAL Ultra

Correction of strongly pronounced wrinkles



• Intradermic (deep skin layer), supraperiosteal

Indications for use:

- Perfect elimination of wrinkles and folds (IV-V degrees)
- Restoration of the volume of soft tissue in the malar, temporal and nasal areas
- Restoration of volume in the chin area, facial contouring
- Volumetric restoration of soft tissue in the buccal, temporal, zygomatic areas



CRYSTAL Ultra

Product Details

PACKAGE	1 x 1.0 ml
HA CONCENTRATION	20 mg/ml
LIDOCAINE	0.3 %
NEEDLES IN ASSORTMENT	25 G ½" - 1 ea. 27 G ½" - 1 ea.
STABILIZATION	BDDE
CROSS-LINK LEVEL	
DURATION*	Up to 18 months

Package

1 box: 1 syringe / 2 needles

* Duration period can depend on the injection method, area of injection and the patient's individual characteristics.



Storage

Avoid direct sunlight and store at 2°C to 25°C

Benefits

Benefits

- Use of highly **purified hyaluronic acid** (non-animal origin)
- No bacterial endotoxins
- **High safety** due to low concentrations of the residual cross-linking agent
- Long lasting effect
- Low risk of scars due to low level of protein left in HA gel
- No migration and no lumps due to highly elastic formula



Treatment areas

















RYSTAL®

wrinkles **Correction of**

Correction of wrinkles



BEFORE



AFTER



BEFORE




AFTER

Volumetric lip modelling

Volumetric lip modelling







Augmentation and lip volumizing

Shape changing

Recreating and rebuilding lip contour

Asymmetry correction

Correction of neck and chin areas

- Correcting aesthetic defects of the face oval
- Reducing flabbiness of the skin under the chin
- Correcting horizontal wrinkles on the neck
- Eliminating initial signs of gravitational ptosis



Face contouring

- Face contour modelling
- Volumetric remodelling of the chin
- Remodelling of the mandibular angles



for the hands

Correction for the back of the hands

- Correcting wrinkles on the back of the hands
- Volumetric modelling of the dorsum of the hands
- Creating optimal amount of soft tissue and improving appearance of the hands



Non-surgical rhinoplasty

Non-surgical rhinoplasty

- Nose bridge correction
- Nose tip correction
- Asymmetry correction
- Hump correction










AFTER



Precautions

Anatomically dangerous areas

- The area of the medial angle of the eye the place of passage a. et v. angularis
- Medial segment of the lacrimal sulcus, 1 cm from the inner canthus
- The place of exit of the supraorbital neurovascular bundle is along the mid-pupillary line of the eyebrow
- The exit point of the infraorbital neurovascular bundle is 0.8 - 1.0 cm below the bone edge of the orbit, at the level of the medial edge of the iris
- The projection area of the cheekbone-facial nerve is 5-7 mm lateral to the inferior arcuate orbit edge
- Place of exit of the mental neurovascular bundle - in the midpupillary line